


University-level worksheet · greyscale print version

Aromatic Chemistry

the island



the positive charge is shared over three carbons (positions 2, 4 and 6 relative to the attacked carbon)

What this is

Seven short sections, one for each lesson of 6.1.1, containing the first-year university explanation of what the lesson taught: why $4n + 2$, what the horseshoe really stands for, why phenol is acidic, why groups direct where they do. None of it is examined at A level, and none of it appears on the ordinary worksheets — it lives here so that nobody meets it by accident.

Who it is for

Anyone who has finished the Core and Harder questions and wants to know why. If you enjoy this, the Stretch Pack (63 pages, first- and second-year level) is the next step; each section here says where to go in it.

Contents

- U1** Aromaticity: why $4n + 2$? (lesson 1)
- U2** How chemists really name rings: ortho, meta, para and trivial names (lesson 2)
- U3** The Wheland intermediate (lesson 3)
- U4** Halogen carriers are Lewis acids (lesson 4)
- U5** Friedel–Crafts: the fine print (lesson 5)
- U6** Phenol: acidity explained properly (lesson 6)
- U7** Directing effects: the real reason (lesson 7)

Worked answers at the back

U1


GOES WITH LESSON 1 · UNIVERSITY LEVEL · NOT EXAMINED

Aromaticity: why $4n + 2$?


Benzene is not the only aromatic ring — and some rings with alternating double bonds are spectacularly UNstable.

Key facts


- Hückel's rule: a planar, fully conjugated ring is aromatic if it holds $4n + 2$ π electrons (2, 6, 10, 14 ...).
- $4n$ π electrons in a planar ring is ANTiaromatic — far less stable than the open-chain version. Cyclobutadiene (4 π) survives only in an argon matrix at 8 K.
- A ring that escapes planarity is merely non-aromatic: cyclooctatetraene (8 π) puckers into a tub and behaves like an alkene.
- Lone pairs count only if they sit in a p-orbital that is part of the ring: pyrrole's N lone pair does; pyridine's does not.




naphthalene



pyridine






pyrrole



tropylium cation

Aromatic rings that are not benzene: naphthalene (10 π), pyridine (6 π), pyrrole (6 π including the N lone pair), tropylium cation (6 π).

benzene: 6 π electrons**cyclobutadiene: 4 π electrons****cyclopentadienyl anion: 6 π** 

Frost circles: the π orbital energies of a ring sit at the corners of the polygon inscribed vertex-down in a circle. $4n + 2$ electrons fill a closed shell.

U1.1 Classify as aromatic, antiaromatic or non-aromatic, giving the π -electron count: (a) the cyclopropenyl cation C_3H_3^+ (b) cyclobutadiene C_4H_4 (c) cyclooctatetraene C_8H_8 (d) the cyclopentadienyl anion C_5H_5^- (e) naphthalene.

U1.2 Naphthalene's C–C bonds are NOT all the same length (0.136–0.142 nm). Explain why it is still aromatic, and why its bonds differ when benzene's do not.

U1.3 Use a Frost circle (square, vertex down) to explain why planar cyclobutadiene would be a diradical.

Go further: Stretch Pack Part A §1 (aromaticity) and Part B §8 (Hückel molecular orbital theory with numbers).

U2

GOES WITH LESSON 2 · UNIVERSITY LEVEL · NOT EXAMINED

How chemists really name rings: ortho, meta, para and trivial names

The names you will meet in every university textbook and paper, and why the exam board avoids them.

Key facts

- ortho (o-) = positions 1,2; meta (m-) = 1,3; para (p-) = 1,4. Only meaningful for TWO groups.
- Trivial names still in daily use: toluene (methylbenzene), aniline (phenylamine), xylene (dimethylbenzene), cresol (methylphenol), anisole (methoxybenzene), benzaldehyde and benzoic acid (also IUPAC), styrene (phenylethene).
- IUPAC lowest-locant and alphabetical rules still apply underneath: p-xylene is 1,4-dimethylbenzene.

U2.1 Give the IUPAC names of: p-xylene, m-cresol, o-nitrotoluene, p-chloroaniline, anisole.

U2.2 Explain why 1,2,4-trichlorobenzene cannot be named with ortho/meta/para, and why 'o-dichlorobenzene' is unambiguous.

U2.3 'Aryl' (Ar-) is used the way 'alkyl' (R-) is. Write the general formula for an aryl chloride and for Friedel–Crafts acylation of an arene, using Ar.


Go further: Stretch Pack Part B §11 (regioselectivity and synthesis) uses o/m/p throughout.

U3

GOES WITH LESSON 3 · UNIVERSITY LEVEL · NOT EXAMINED


The Wheland intermediate

The horseshoe is a shorthand. Here is what it stands for, and why the intermediate loses H^+ rather than adding a nucleophile.



the positive charge is shared over three carbons (positions 2, 4 and 6 relative to the attacked carbon)

The three resonance forms of the intermediate (arenium ion, σ -complex): the positive charge is shared over the carbons ortho and para to the attacked carbon.



Energy profile: step 1 (attack, aromaticity lost) is slow and rate-determining; step 2 (loss of H^+ , aromaticity regained) is fast.

Key facts

- The intermediate is real: heptamethylbenzenium salts have been isolated and characterised by NMR (Doering 1958; Olah in superacids).
- Nitration of C_6D_6 is as fast as C_6H_6 — no kinetic isotope effect — so the C–H bond is not broken in the rate-determining step (Melander 1950).
- Substitution beats addition because losing H^+ regains $\sim 150 \text{ kJ mol}^{-1}$ of delocalisation energy; addition would give a non-aromatic diene.

U3.1 Draw the three resonance forms of the intermediate formed when Br^+ attacks benzene. Mark the carbon that carries H and Br, and state on which carbons the positive charge is found.

U3.2 Use the energy profile to explain why the intermediate loses H^+ rather than picking up HSO_4^- (which would give an addition product, as happens with an alkene).

U3.3 Sulfonation (SO_3) shows a kinetic isotope effect and is reversible; nitration shows neither. Sketch the two energy profiles and explain the difference.

U3.4 NO_2^+ is linear. Which common molecule is it isoelectronic with, and what does linearity tell you about the N atom's orbitals?

Go further: Stretch Pack Part A §3 (the Wheland intermediate) and Part B §10 (electrophilic substitution at research depth).

U4

GOES WITH LESSON 4 · UNIVERSITY LEVEL · NOT EXAMINED

Halogen carriers are Lewis acids*What AlCl_3 and FeBr_3 actually do, and why iodination is hard.***Key facts**

- A Lewis acid is an electron-pair acceptor. AlCl_3 has only six electrons around Al; FeBr_3 has empty d/p orbitals on Fe.
- The real electrophile is often a polarised complex $\text{Br}-\text{Br}\cdots\text{FeBr}_3$ ($\text{Br}\delta^+$) rather than free Br^+ ; the exam board lets you write Br^+ because the arrow-pushing is identical.
- Iron filings work because $2\text{Fe} + 3\text{Br}_2 \rightarrow 2\text{FeBr}_3$ forms the carrier in situ.

U4.1 Draw a curly-arrow mechanism for the formation of the electrophile from Br_2 and FeBr_3 , showing (a) the $\text{Br}-\text{Br}$ bond pair moving to Fe and (b) the resulting FeBr_4^- and Br^+ .

U4.2 Iodobenzene cannot be made efficiently from benzene, I_2 and FeI_3 . Give two reasons, and suggest why adding nitric acid helps.

U4.3 Fluorination of benzene with F_2 is violently exothermic and gives a tar. Explain why direct fluorination is useless for making fluorobenzene, and name the general strategy chemists use instead.

Go further: Stretch Pack Part A §5 (Friedel–Crafts fine print) and Part B §11 (diazonium chemistry).

U5

GOES WITH LESSON 5 · UNIVERSITY LEVEL · NOT EXAMINED

Friedel–Crafts: the fine print*Four things the A-level story leaves out — and the standard fix for all of them.***Key facts**

- Primary carbocations rearrange to more stable secondary/tertiary ones by hydride or methyl shifts before they attack the ring.
- Alkyl groups activate the ring, so alkylation often goes further (polyalkylation) unless benzene is in large excess.
- Strongly deactivated rings ($-\text{NO}_2$, $-\text{COR}$, $-\text{CN}$) do not undergo Friedel–Crafts reactions at all; $-\text{NH}_2$ and $-\text{OH}$ poison the catalyst by complexing AlCl_3 .
- Acylation avoids all of this: the acylium ion is resonance-stabilised and does not rearrange, and the ketone product is deactivated so the reaction stops after one substitution.

U5.1 Benzene + 1-chloropropane/ AlCl_3 gives mainly (1-methylethyl)benzene. Draw the carbocation rearrangement responsible and explain the driving force.

U5.2 Design a two-step synthesis of propylbenzene from benzene that avoids rearrangement, naming the reagents for each step.

U5.3 Draw the two resonance forms of the ethanoyl cation and explain why it does not rearrange.

U5.4 Why is nitrobenzene a good SOLVENT for Friedel–Crafts acylations, whereas methoxybenzene is not?


Go further: Stretch Pack Part A §5 and Part B §12 (named reactions and rearrangements).

U6


GOES WITH LESSON 6 · UNIVERSITY LEVEL · NOT EXAMINED

Phenol: acidity explained properly

Why phenol is 10^6 times more acidic than ethanol yet 10^5 times less than ethanoic acid — and what a nitro group does.



phenoxide ion



negative charge on a ring carbon
(one of three such forms: positions 2, 4, 6)

The phenoxide ion: the negative charge on oxygen is delocalised onto C2, C4 and C6 (three extra resonance forms).

Key facts

- pK_a : ethanol ≈ 16 ; phenol 10.0; 4-nitrophenol 7.2; 2,4-dinitrophenol 4.1; 2,4,6-trinitrophenol (picric acid) 0.4; ethanoic acid 4.8.
- An acid is stronger when its conjugate base is more stable: delocalisation of the negative charge stabilises the phenoxide but not the ethoxide ion.
- Nitro groups withdraw electron density inductively ($-I$) AND accept the negative charge by resonance when para or ortho ($-M$), so each one strengthens the acid dramatically.

U6.1 Explain, using the resonance forms above, why phenol is a much stronger acid than ethanol.

U6.2 Explain the trend phenol (10.0) → 4-nitrophenol (7.2) → picric acid (0.4). Why is 3-nitrophenol (8.4) a weaker acid than 4-nitrophenol?

U6.3 Aspirin is made from 2-hydroxybenzoic acid, itself made from phenol by the Kolbe–Schmitt reaction (sodium phenoxide + CO₂ at 125 °C and 100 atm, then acid). Suggest why the phenoxide ion, not phenol, is used, and why the CO₂ goes to the 2-position.


Go further: Stretch Pack Part A §6 (phenol acidity, Kolbe–Schmitt, aspirin).

U7

GOES WITH LESSON 7 · UNIVERSITY LEVEL · NOT EXAMINED

Directing effects: the real reason

Count the resonance forms of the intermediate and the 2,4- versus 3- rule explains itself — and partial rate factors let you predict the isomer ratio.



attack at position 2 (ortho): the oxygen lone pair gives a FOURTH resonance form — every atom has a full octet

Attack at the 2-position of phenol: the fourth resonance form, in which the oxygen lone pair makes a $C=O^+$ bond, gives every atom a full octet — attack at 3 cannot do this.

Key facts

- For $-OH$, $-NH_2$, $-OR$, $-Cl$: attack at 2 or 4 gives an EXTRA resonance form stabilised by the heteroatom lone pair; attack at 3 does not.
- For $-NO_2$, $-COR$: attack at 2 or 4 puts the positive charge next to the δ^+ atom of the group (very unfavourable); attack at 3 avoids this — so every position is deactivated, but 3 least.
- Partial rate factors compare ONE position of the substituted ring with ONE position of benzene. Nitration of methylbenzene: ortho 42, meta 2.5, para 58. Chlorobenzene: 0.030, 0.0009, 0.14.

U7.1 Use the methylbenzene partial rate factors to calculate the expected percentages of 2-, 3- and 4-nitromethylbenzene, and the overall rate relative to benzene.

U7.2 Every partial rate factor for methylbenzene is greater than 1, even the meta one. What does this tell you about the $-CH_3$ group?

U7.3 Explain why $-\text{Cl}$ deactivates the ring (all partial rate factors below 1) yet directs to 2 and 4.

U7.4 Draw the three resonance forms for attack of NO_2^+ at the 4-position of nitrobenzene and identify the one that makes 4-attack unfavourable.

Go further: Stretch Pack Part A §4 (directing effects) and Part B §9 (the Hammett equation — putting numbers on 'electron-withdrawing').

Worked answers

Argue with these. If your answer is different and you can defend it, bring it to Dr Fennell.

U1 — Aromaticity: why $4n + 2$?

U1.1 Classify as aromatic, antiaromatic or non-aromatic, giving the π -electron count: (a) the cyclopropenyl cation $C_3H_3^+$ (b) cyclobutadiene...

(a) 2π ($n = 0$): aromatic — the smallest aromatic ring. (b) $4\pi = 4n$: antiaromatic if planar; it distorts to a rectangle and is extremely reactive. (c) 8π : would be antiaromatic if planar, so it adopts a tub shape — non-aromatic, alkene-like. (d) 6π (4 from $C=C$ + the lone pair): aromatic — which is why cyclopentadiene ($pK_a \approx 16$) is a remarkably strong acid for a hydrocarbon. (e) 10π ($n = 2$): aromatic.

U1.2 Naphthalene's C–C bonds are NOT all the same length (0.136–0.142 nm). Explain why it is still aromatic, and why its bonds differ w...

Aromaticity needs a closed shell of $4n + 2$ delocalised π electrons, which naphthalene has (10). Bond lengths depend on how often each bond is double in the resonance structures: naphthalene has three Kekulé structures and C1–C2 is double in two of them (bond order 1.67, 0.136 nm) while C2–C3 is double in only one (1.33, 0.142 nm). In benzene every bond is double in one of the two structures, so all are identical (order 1.5).

U1.3 Use a Frost circle (square, vertex down) to explain why planar cyclobutadiene would be a diradical.

The four orbital energies sit at the corners of the square: one bonding (bottom), two degenerate non-bonding (middle), one antibonding (top). Four electrons: two fill the bonding orbital and the remaining two go one each into the degenerate pair (Hund's rule) — two unpaired electrons, no closed shell, no aromatic stabilisation.

U2 — How chemists really name rings: ortho, meta, para and trivial names

U2.1 Give the IUPAC names of: *p*-xylene, *m*-cresol, *o*-nitrotoluene, *p*-chloroaniline, anisole.

1,4-dimethylbenzene; 3-methylphenol; 2-nitromethylbenzene; 4-chlorophenylamine; methoxybenzene.

U2.2 Explain why 1,2,4-trichlorobenzene cannot be named with ortho/meta/para, and why 'o-dichlorobenzene' is unambiguous.

o/*m*/*p* describe the relationship between exactly two substituents; with three groups every pair has its own relationship (1,2 ortho; 1,4 para; 2,4 meta) so the description is ambiguous — locants are required. With two identical groups, 'ortho' fixes the 1,2 relationship completely.

U2.3 'Aryl' (Ar–) is used the way 'alkyl' (R–) is. Write the general formula for an aryl chloride and for Friedel–Crafts acylation of a...

Aryl chloride: $ArCl$. Acylation: $ArH + RCOCl \rightarrow ArCOR + HCl$ ($AlCl_3$ catalyst).

U3 — The Wheland intermediate

U3.1 Draw the three resonance forms of the intermediate formed when Br^+ attacks benzene. Mark the carbon that carries H and Br, and sta...

Ring with H and Br on the same carbon (C1). Positive charge on C2 with $C=C$ at C3–C4 and C5–C6; on C4 with $C=C$ at C2–C3 and C5–C6; on C6 with $C=C$ at C2–C3 and C4–C5. The charge is on the carbons ortho and para to C1 — never on C1 itself and never on C3 or C5. The horseshoe with + is the average of these three drawings.

U3.2 Use the energy profile to explain why the intermediate loses H^+ rather than picking up HSO_4^- (which would give an addition product...)

Loss of H^+ restores the aromatic π -system and releases the delocalisation energy ($\sim 150 \text{ kJ mol}^{-1}$) — a very favourable, low-barrier step. Addition of HSO_4^- would give a non-aromatic cyclohexadiene far higher in energy. An alkene carbocation has no aromaticity to regain, so addition is its only option.

U3.3 Sulfonation (SO_3) shows a kinetic isotope effect and is reversible; nitration shows neither. Sketch the two energy profiles and ex...

Sulfonation: the two transition states are of similar height because SO_3 is a weaker electrophile and the intermediate can revert to reactants; C–H (C–D) cleavage is partly rate-determining, so C_6D_6 reacts more slowly, and the shallow well makes the reaction reversible (steam removes $-SO_3H$). Nitration: the first transition state is much higher than the second, so the C–H step never limits the rate and the reaction is effectively irreversible.

U3.4 NO_2^+ is linear. Which common molecule is it isoelectronic with, and what does linearity tell you about the N atom's orbitals?

CO_2 (16 valence electrons, $O=N^+=O$). Linear $\rightarrow$ the N is sp hybridised with two π bonds; the empty p-orbital character at N is what makes it such a powerful electrophile.

U4 — Halogen carriers are Lewis acids

U4.1 Draw a curly-arrow mechanism for the formation of the electrophile from Br_2 and $FeBr_3$, showing (a) the Br–Br bond pair moving to F...

Arrow from a lone pair on one Br to the empty orbital on Fe gives $Br-Br \rightarrow FeBr_3$; then an arrow from the Br–Br bond onto the Fe-bound Br gives $FeBr_4^-$ and Br^+ (in practice the ring attacks the polarised complex directly).

U4.2 Iodobenzene cannot be made efficiently from benzene, I_2 and FeI_3 . Give two reasons, and suggest why adding nitric acid helps.

I^+ is a poor electrophile (large, polarisable, low charge density), and the reaction is close to thermoneutral/reversible: the HI produced reduces iodobenzene back to benzene. Nitric acid oxidises the HI (and I_2 to a better electrophile, I^+/IO^+ species), driving the reaction forward.

U4.3 Fluorination of benzene with F_2 is violently exothermic and gives a tar. Explain why direct fluorination is useless for making flu...

F_2 is such a strong oxidant and the C–F bond so strong that the reaction is uncontrolled (radical pathways, polysubstitution, ring destruction). Fluoroarenes are made indirectly — e.g. from the amine via a diazonium salt (Balz–Schiemann: $ArN_2^+BF_4^- \rightarrow ArF + N_2 + BF_3$).

U5 — Friedel–Crafts: the fine print

U5.1 Benzene + 1-chloropropane/ $AlCl_3$ gives mainly (1-methylethyl)benzene. Draw the carbocation rearrangement responsible and explain th...

$CH_3CH_2CH_2^+$ (primary) undergoes a 1,2-hydride shift: a hydrogen with its bonding pair moves from C2 to C1, giving $CH_3CH^+CH_3$ (secondary). Secondary cations are more stable because two alkyl groups donate electron density inductively and by hyperconjugation.

U5.2 Design a two-step synthesis of propylbenzene from benzene that avoids rearrangement, naming the reagents for each step.

1 Acylate with $CH_3CH_2COCl/AlCl_3 \rightarrow$ 1-phenylpropan-1-one (acylium ions do not rearrange; the product is deactivated so no second acylation). 2 Reduce the C=O to CH_2 : Clemmensen ($Zn(Hg)$, conc. HCl, heat) or Wolff–Kishner (N_2H_4 , KOH, heat).

U5.3 Draw the two resonance forms of the ethanoyl cation and explain why it does not rearrange.

$\text{CH}_3\text{—C}\equiv\text{O}^+ \leftrightarrow \text{CH}_3\text{—C}^+=\text{O}$. The positive charge is delocalised onto oxygen (every atom keeps an octet in the first form), so the cation is far more stable than an alkyl cation and has no driving force to rearrange.

U5.4 Why is nitrobenzene a good SOLVENT for Friedel–Crafts acylations, whereas methoxybenzene is not?

Nitrobenzene is so deactivated by —NO_2 that it does not react with acylium ions, yet it dissolves AlCl_3 complexes well. Methoxybenzene is activated (+M) and would be acylated itself, at the para position, faster than benzene.

U6 — Phenol: acidity explained properly

U6.1 Explain, using the resonance forms above, why phenol is a much stronger acid than ethanol.

In the phenoxide ion the negative charge is delocalised from O onto the ring carbons (three resonance forms with the charge on C2, C4 and C6), which stabilises the anion and shifts the dissociation equilibrium to the right. The ethoxide ion has no delocalisation — the full charge stays on one oxygen.

U6.2 Explain the trend phenol (10.0) → 4-nitrophenol (7.2) → picric acid (0.4). Why is 3-nitrophenol (8.4) a weaker acid than 4-nitroph...

Each —NO_2 withdraws electron density (–I) and, from the 2- or 4-position, accepts the negative charge by resonance (an extra form with the charge on a nitro oxygen), stabilising the anion further; three nitro groups make picric acid as strong as a mineral acid. From the 3-position the charge cannot be pushed onto the nitro group (no resonance form reaches it), so only the weaker inductive effect operates.

U6.3 Aspirin is made from 2-hydroxybenzoic acid, itself made from phenol by the Kolbe–Schmitt reaction (sodium phenoxide + CO_2 at 125 °...

O^- donates far more electron density into the ring than OH, activating it enough to attack the weak electrophile CO_2 . The Na^+ ion coordinates both the phenoxide oxygen and CO_2 , delivering the electrophile to the adjacent (2-) carbon.

U7 — Directing effects: the real reason

U7.1 Use the methylbenzene partial rate factors to calculate the expected percentages of 2-, 3- and 4-nitromethylbenzene, and the overa...

Weight each position by the number of equivalent positions: ortho $2 \times 42 = 84$; meta $2 \times 2.5 = 5$; para 58; total 147. 2-: $84/147 = 57\%$; 3-: 3.4% ; 4-: 39% (experiment $\approx 58 : 4 : 38$). Overall rate = $147/6 \approx 25$ times faster than benzene.

U7.2 Every partial rate factor for methylbenzene is greater than 1, even the meta one. What does this tell you about the —CH_3 group?

—CH_3 activates ALL positions (inductive electron donation raises the electron density everywhere) but ortho/para far more, because hyperconjugation stabilises the intermediates for attack at those positions.

U7.3 Explain why —Cl deactivates the ring (all partial rate factors below 1) yet directs to 2 and 4.

Inductive withdrawal (Cl is electronegative) lowers the electron density at every position, so chlorobenzene reacts more slowly than benzene. But a Cl lone pair can donate by resonance into the intermediates for ortho/para attack (an extra resonance form), so those positions are deactivated less than meta → 2,4-directing.

U7.4 Draw the three resonance forms for attack of NO_2^+ at the 4-position of nitrobenzene and identify the one that makes 4-attack unfav...

One of the three forms puts the positive charge on the carbon bearing $-\text{NO}_2$, directly next to the δ^+ nitrogen — like charges adjacent, and no lone pair to help. This form contributes little, so the intermediate is less stabilised than for 3-attack, where the charge never reaches that carbon.